



TRANSLATOR'S DECLARATION

I, Janet Hope, BSc (Hons.), MIL., MITI., translator to Messrs. Taylor and Meyer of 20 Kingsmead Road, London, SW2 3JD, Great Britain, verify that I know well both the German and the English language, that I have prepared the attached English translation of 33 pages of a German Patent application in the German language with the title:

Neue für die Gene metR und metZ kodierende Nukleotidsequenzen

identified by the code number 000369 BT at the upper left of each page and corresponding to client/matter number of the law firm of

and that the attached English translation of this document is a true and correct translation of the document attached thereto to the best of my knowledge and belief.

I further declare that all statements made of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements are made with the knowledge that wilful false statements and the like are punishable by fine or imprisonment, or both, under 18 USC 1001, and that such false statements may jeopardize the validity of this document.

By: _____

Date: 3rd October 2003

FEDERAL REPUBLIC OF GERMANY

Certificate of Priority for Filing of a Patent Application

Filing number: 100 43 335.9

Filing date: 2nd September 2000

Applicant/Proprietor: Degussa AG, Düsseldorf/Germany

First applicant: Degussa-Hüls
Aktiengesellschaft, Frankfurt-am-
Main/Germany

Title: New nucleotide sequences which code
for the metR and metZ genes

IPC: C 12 N, C 12 Q, C 07 H

The attached papers are a true and accurate reproduction of
the original documents for this patent application.

Munich, 13th June 2001
On behalf of the President
of the German Patent and Trade
Mark Office

(signature)

Faust

**New nucleotide sequences which code for the metR
and metZ genes**

The invention provides nucleotide sequences from coryneform bacteria which code for the metR and metZ genes and a
5 process for the fermentative preparation of amino acids, in particular L-methionine, by attenuation of the metR and/or metZ gene.

Prior art

10 L-Amino acids, in particular methionine, are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly in animal nutrition.

It is known that amino acids are prepared by fermentation from strains of coryneform bacteria, in particular
15 Corynebacterium glutamicum. Because of their great importance, work is constantly being undertaken to improve the preparation processes. Improvements to the process can relate to fermentation measures, such as, for example, stirring and supply of oxygen, or the composition of the
20 nutrient media, such as, for example, the sugar concentration during the fermentation, or the working up to the product form by, for example, ion exchange chromatography, or the intrinsic output properties of the microorganism itself.

25 Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites or are auxotrophic for metabolites of regulatory importance and which produce amino acids, such
30 as e.g. L-methionine, are obtained in this manner.

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of Corynebacterium strains which produce L-amino acids, by

amplifying individual amino acid biosynthesis genes and investigating the effect on the amino acid production.

Object of the invention

The inventors had the object of providing new measures for improved fermentative preparation of amino acids, in particular L-methionine.

Description of the invention

Where L-amino acids or amino acids are mentioned in the following, this means one or more amino acids, including their salts, chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine.

When L-methionine or methionine are mentioned in the following, the salts, such as e.g. methionine hydrochloride or methionine sulfate are also meant by this.

The invention provides isolated polynucleotides from coryneform bacteria, which comprise the polynucleotide sequences which code for the metR and/or metZ genes, chosen from the group consisting of

a) polynucleotide which is identical to the extent of at least 70% to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 2,

b) polynucleotide which is identical to the extent of at least 70% to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 3,

c) polynucleotide which codes for a polypeptide which comprises an amino acid sequence which is identical to

the extent of at least 70% to the amino acid sequence of
SEQ ID No. 2,

- d) polynucleotide which codes for a polypeptide which
comprises an amino acid sequence which is identical to
5 the extent of at least 70% to the amino acid sequence of
SEQ ID No. 3,
- e) polynucleotide which is complementary to the
polynucleotides of a), b), c) or d), and
- f) polynucleotide comprising at least 15 successive
10 nucleotides of the polynucleotide sequence of a), b),
c), d) or e),

the polypeptides according to a) or c) preferably having
the activity of the transcription activator MetR and the
polypeptides according to b) or d) preferably having the
15 activity of O-succinylhomoserine sulfhydrylase (MetZ).

The invention also provides the abovementioned
polynucleotides, these preferably being DNAs which are
capable of replication, comprising:

- (i) the nucleotide sequence shown in SEQ ID No. 1, or
- 20 (ii) at least one sequence which corresponds to
sequence (i) within the range of the degeneration
of the genetic code, or
- (iii) at least one sequence which hybridizes with the
sequences complementary to sequences (i) or (ii),
25 and optionally
- (iv) sense mutations of neutral function in (i).

The invention also provides:

a DNA which is capable of replication and comprises the
nucleotide sequence as shown in SEQ ID No. 1,

a polynucleotide which codes for a polypeptide which comprises the amino acid sequence as shown in SEQ ID No. 2 or SEQ ID No. 3,

5 a vector containing parts of the polynucleotide according to the invention, but at least 15 successive nucleotides of the sequence claimed

and coryneform bacteria in which the metR gene and/or the metZ gene is or are attenuated, in particular by deletion, insertion or base exchange.

10 The invention also provides polynucleotides which substantially comprise a polynucleotide sequence, which are obtainable by screening by means of hybridization of a corresponding gene library of a coryneform bacterium, which comprises the complete gene or parts thereof, with a probe
15 which comprises the sequence of the polynucleotide according to the invention according to SEQ ID No. 1 or a fragment thereof, and isolation of the polynucleotide sequence mentioned.

Polynucleotides which comprise the sequences according to
20 the invention are suitable as hybridization probes for RNA, cDNA and DNA, in order to isolate, in the full length, nucleic acids, or polynucleotides or genes which code for the transcription activator MetR and/or O-succinylhomoserine sulfhydrylase or to isolate those
25 nucleic acids or polynucleotides or genes which have a high similarity with the sequence of the transcription activator MetR gene and/or that of the O-succinylhomoserine sulfhydrylase gene.

Polynucleotides which comprise the sequences according to
30 the invention are furthermore suitable as primers with the aid of which DNA of genes which code for the transcription activator MetR and/or O-succinylhomoserine sulfhydrylase can be prepared by the polymerase chain reaction (PCR).

Such oligonucleotides which serve as probes or primers comprise at least 30, preferably at least 20, very particularly preferably at least 15 successive nucleotides. Oligonucleotides which have a length of at least 40 or 50 nucleotides are also suitable.

"Isolated" means separated out of its natural environment.

"Polynucleotide" in general relates to polyribonucleotides and polydeoxyribonucleotides, it being possible for these to be non-modified RNA or DNA or modified RNA or DNA.

10 The polynucleotides according to the invention include a polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom and also those which are at least 70%, preferably at least 80% and in particular at least 90% to 95% identical to the polynucleotide according to SEQ ID
15 No. 1 or a fragment prepared therefrom.

The polypeptides according to the invention include the polypeptides according to SEQ ID No. 2 and SEQ ID No. 3, in particular those with the biological activity of the transcription activator MetR and of O-succinylhomoserine
20 sulfhydrylase, and also those which are at least 70%, preferably at least 80%, and in particular which are at least 90% to 95% identical to the polypeptides according to SEQ ID No. 2 and SEQ ID No. 3 and have the activities mentioned.

25 "Polypeptides" are understood as meaning peptides or proteins which comprise two or more amino acids bonded via peptide bonds.

The invention moreover provides a process for the fermentative preparation of amino acids, in particular
30 methionine, using coryneform bacteria which in particular already produce the amino acids, and in which the nucleotide sequences which code for the metR gene and/or

for the metZ gene are attenuated, in particular eliminated or expressed at a low level.

The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by using a weak promoter or using a gene or allele which codes for a corresponding enzyme with a low activity or inactivates the corresponding gene or enzyme (protein), and optionally combining these measures.

The microorganisms which the present invention provides can prepare L-amino acids, in particular methionine, from glucose, sucrose, lactose, fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They can be representatives of coryneform bacteria, in particular of the genus *Corynebacterium*. Of the genus *Corynebacterium*, there may be mentioned in particular the species *Corynebacterium glutamicum*, which is known among experts for its ability to produce L-amino acids.

Suitable strains of the genus *Corynebacterium*, in particular of the species *Corynebacterium glutamicum* (*C. glutamicum*), are in particular the known wild-type strains

Corynebacterium glutamicum ATCC13032
Corynebacterium acetoglutamicum ATCC15806
Corynebacterium acetoacidophilum ATCC13870
Corynebacterium melassecola ATCC17965
Corynebacterium thermoaminogenes FERM BP-1539
Brevibacterium flavum ATCC14067
Brevibacterium lactofermentum ATCC13869 and
Brevibacterium divaricatum ATCC14020

or L-amino acid-producing mutants or strains prepared therefrom, such as, for example, the L-methionine-producing strain

Corynebacterium glutamicum ATCC21608.

The inventors have succeeded in isolating the new metR and metZ genes from *C. glutamicum* which code for the transcription activator MetR and the enzyme O-succinylhomoserine sulfhydrylase.

To isolate the metR gene, the metZ gene or also other genes of *C. glutamicum*, a gene library of this microorganism is first set up in *Escherichia coli* (*E. coli*). The setting up of gene libraries is described in generally known textbooks and handbooks. The textbook by Winnacker: Gene und Klone, Eine Einführung in die Gentechnologie (Verlag Chemie, Weinheim, Germany, 1990), or the handbook by Sambrook et al.: Molecular Cloning, A Laboratory Manual (Cold Spring Harbor Laboratory Press, 1989) may be mentioned as an example. A well-known gene library is that of the *E. coli* K-12 strain W3110 set up in λ vectors by Kohara et al. (Cell 50, 495-508 (1987)). Bathe et al. (Molecular and General Genetics, 252:255-265, 1996) describe a gene library of *C. glutamicum* ATCC13032, which was set up with the aid of the cosmid vector SuperCos I (Wahl et al., 1987, Proceedings of the National Academy of Sciences USA, 84:2160-2164) in the *E. coli* K-12 strain NM554 (Raleigh et al., 1988, Nucleic Acids Research 16:1563-1575).

Börmann et al. (Molecular Microbiology 6(3), 317-326 (1992)) in turn describe a gene library of *C. glutamicum* ATCC13032 using the cosmid pHC79 (Hohn and Collins, Gene 11, 291-298 (1980)). To prepare a gene library of *C. glutamicum* in *E. coli* it is also possible to use plasmids such as pBR322 (Bolivar, Life Sciences, 25, 807-818 (1979)) or pUC9 (Vieira et al., 1982, Gene, 19:259-268). Suitable hosts are, in particular, those *E. coli* strains which are restriction- and recombination-defective. An example of these is the strain DH5 α mcr, which has been described by Grant et al. (Proceedings of the National Academy of Sciences USA, 87 (1990) 4645-4649). The long DNA fragments

cloned with the aid of cosmids can in turn be subcloned in the usual vectors suitable for sequencing and then sequenced, as is described e.g. by Sanger et al.

(Proceedings of the National Academy of Sciences of the United States of America, 74:5463-5467, 1977).

The resulting DNA sequences can then be investigated with known algorithms or sequence analysis programs, such as e.g. that of Staden (Nucleic Acids Research 14, 217-232 (1986)), that of Marck (Nucleic Acids Research 16, 1829-1836 (1988)) or the GCG program of Butler (Methods of Biochemical Analysis 39, 74-97 (1998)).

The new DNA sequence of *C. glutamicum* which codes for the metR and metZ genes and which, as SEQ ID No. 1, is a constituent of the present invention has been found in this manner. The amino acid sequence of the corresponding proteins has furthermore been derived from the present DNA sequence by the methods described above. The resulting amino acid sequences of the metR and metZ gene products are shown in SEQ ID No. 2 and SEQ ID No. 3.

Coding DNA sequences which result from SEQ ID No. 1 by the degeneracy of the genetic code are also a constituent of the invention. Conservative amino acid exchanges, such as e.g. exchange of glycine for alanine or of aspartic acid for glutamic acid in proteins, are furthermore known among experts as "sense mutations" which do not lead to a fundamental change in the activity of the protein, i.e. are of neutral function. It is furthermore known that changes on the N and/or C terminus of a protein cannot substantially impair or can even stabilize the function thereof. Information in this context can be found by the expert, inter alia, in Ben-Bassat et al. (Journal of Bacteriology 169:751-757 (1987)), in O'Regan et al. (Gene 77:237-251 (1989)), in Sahin-Toth et al. (Protein Sciences 3:240-247 (1994)), in Hochuli et al. (Bio/Technology 6:1321-1325 (1988)) and in known textbooks of genetics and

molecular biology. Amino acid sequences which result in a corresponding manner from SEQ ID No. 2 or SEQ ID No. 3 are also a constituent of the invention.

In the same way, DNA sequences which hybridize with SEQ ID
5 No. 1 or parts of SEQ ID No. 1 are a constituent of the invention. Finally, DNA sequences which are prepared by the polymerase chain reaction (PCR) using primers which result from SEQ ID No. 1 are a constituent of the invention. Such oligonucleotides typically have a length
10 of at least 15 nucleotides.

Instructions for identifying DNA sequences by means of hybridization can be found by the expert, inter alia, in the handbook "The DIG System Users Guide for Filter Hybridization" from Boehringer Mannheim GmbH (Mannheim,
15 Germany, 1993) and in Liebl et al. (International Journal of Systematic Bacteriology (1991) 41: 255-260). The hybridization takes place under stringent conditions, i.e. only hybrids in which the probe and target sequence, i.e. the polynucleotides treated with the probe, are at least
20 70% identical are formed. It is known that the stringency of the hybridization, including the washing steps, is influenced or determined by varying the buffer composition, the temperature and the salt concentration. The hybridization reaction is preferably carried out under a
25 relatively low stringency compared with the washing steps (Hybaid Hybridisation Guide, Hybaid Limited, Teddington, UK, 1996).

A 5x SSC buffer at a temperature of approx. 50 - 68°C, for example, can be employed for the hybridization reaction.
30 Probes can also hybridize here with polynucleotides which are less than 70% identical to the sequence of the probe. Such hybrids are less stable and are removed by washing under stringent conditions. This can be achieved, for example, by lowering the salt concentration to 2x SSC and
35 optionally subsequently 0.5x SSC (The DIG System User's

Guide for Filter Hybridisation, Boehringer Mannheim, Mannheim, Germany, 1995) a temperature of approx. 50 - 68°C being established. It is optionally possible to lower the salt concentration to 0.1x SSC. Polynucleotide fragments which are, for example, at least 70% or at least 80% or at least 90% to 95% identical to the sequence of the probe employed can be isolated by increasing the hybridization temperature stepwise from 50 to 68°C in steps of approx. 1 - 2°C. Further instructions on hybridization are obtainable on the market in the form of so-called kits (e.g. DIG Easy Hyb from Roche Diagnostics GmbH, Mannheim, Germany, Catalogue No. 1603558).

Instructions for amplification of DNA sequences with the aid of the polymerase chain reaction (PCR) can be found by the expert, inter alia, in the handbook by Gait: Oligonucleotide synthesis: A Practical Approach (IRL Press, Oxford, UK, 1984) and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994).

During work on the present invention, it was found that coryneform bacteria produce amino acids, in particular L-methionine, in an improved manner after attenuation of the metR and/or metZ gene.

To achieve an attenuation, either the expression of the metR and/or of the metZ gene or the catalytic properties of the enzyme proteins can be reduced or eliminated. The two measures can optionally be combined.

The reduction in gene expression can take place by suitable culturing or by genetic modification (mutation) of the signal structures of gene expression. Signal structures of gene expression are, for example, repressor genes, activator genes, operators, promoters, attenuators, ribosome binding sites, the start codon and terminators. The expert can find information on this e.g. in the patent application WO 96/15246, in Boyd and Murphy (Journal of Bacteriology 170: 5949 (1988)), in Voskuil and Chambliss

- (Nucleic Acids Research 26: 3548 (1998), in Jensen and Hammer (Biotechnology and Bioengineering 58: 191 (1998)), in Pátek et al. (Microbiology 142: 1297 (1996)), Vasicova et al. (Journal of Bacteriology 181: 6188 (1999)) and in
- 5 known textbooks of genetics and molecular biology, such as e.g. the textbook by Knippers ("Molekulare Genetik", 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995) or that by Winnacker ("Gene und Klone", VCH Verlagsgesellschaft, Weinheim, Germany, 1990).
- 10 Mutations which lead to a change or reduction in the catalytic properties of enzyme proteins are known from the prior art; examples which may be mentioned are the works by Qiu and Goodman (Journal of Biological Chemistry 272: 8611-8617 (1997)), Sugimoto et al. (Bioscience Biotechnology and
- 15 Biochemistry 61: 1760-1762 (1997)) and Möckel ("Die Threonindehydratase aus Corynebacterium glutamicum: Aufhebung der allosterischen Regulation und Struktur des Enzyms", Reports from the Jülich Research Centre, Jül-2906, ISSN09442952, Jülich, Germany, 1994). Summarizing
- 20 descriptions can be found in known textbooks of genetics and molecular biology, such as e.g. that by Hagemann ("Allgemeine Genetik", Gustav Fischer Verlag, Stuttgart, 1986).
- Possible mutations are transitions, transversions,
- 25 insertions and deletions. Depending on the effect of the amino acid exchange on the enzyme activity, "missense mutations" or "nonsense mutations" are referred to. Insertions or deletions of at least one base pair in a gene lead to "frame shift mutations", as a consequence of which
- 30 incorrect amino acids are incorporated or translation is interrupted prematurely. Deletions of several codons typically lead to a complete loss of the enzyme activity. Instructions on generation of such mutations are prior art and can be found in known textbooks of genetics and
- 35 molecular biology, such as e.g. the textbook by Knippers ("Molekulare Genetik", 7th edition, Georg Thieme Verlag,

Stuttgart, Germany, 1997), that by Winnacker ("Gene und Klone", VCH Verlagsgesellschaft, Weinheim, Germany, 1999) or that by Hagemann ("Allgemeine Genetik", 4th edition, Gustav Fischer Verlag, Stuttgart, 1999).

- 5 A common method of mutating genes of *C. glutamicum* is the method of "gene disruption" and "gene replacement" described by Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)).

In the method of gene disruption a central part of the
10 coding region of the gene of interest is cloned in a plasmid vector which can replicate in a host (typically *E. coli*), but not in *C. glutamicum*. Possible vectors are, for example, pSUP301 (Simon et al., Bio/Technology 1, 784-791 (1983)), pK18mob or pK19mob (Schäfer et al., Gene 145, 69-
15 73 (1994)), pK18mobsacB or pK19mobsacB (Jäger et al., Journal of Bacteriology 174: 5462-65 (1992)), pGEM-T (Promega corporation, Madison, WI, USA), pCR2.1-TOPO (Shuman (1994). Journal of Biological Chemistry 269:32678-84; US Patent 5,487,993), pCR®Blunt (Invitrogen, Groningen,
20 The Netherlands; Bernard et al., Journal of Molecular Biology, 234: 534-541 (1993)) or pEM1 (Schrumpf et al., 1991, Journal of Bacteriology 173:4510-4516). The plasmid vector which contains the central part of the coding region of the gene is then transferred into the desired strain of
25 *C. glutamicum* by conjugation or transformation. The method of conjugation is described, for example, by Schäfer et al. (Applied and Environmental Microbiology 60, 756-759 (1994)). Methods for transformation are described, for example, by Thierbach et al. (Applied Microbiology and
30 Biotechnology 29, 356-362 (1988)), Dunican and Shivnan (Bio/Technology 7, 1067-1070 (1989)) and Tauch et al. (FEMS Microbiological Letters 123, 343-347 (1994)). After homologous recombination by means of a "cross over" event, the coding region of the gene in question is interrupted by
35 the vector sequence and two incomplete alleles are obtained, one lacking the 3' end and one lacking the 5'

end. This method has been used, for example, by Fitzpatrick et al. (Applied Microbiology and Biotechnology 42, 575-580 (1994)) to eliminate the recA gene of *C. glutamicum*.

5 In the method of "gene replacement", a mutation, such as e.g. a deletion, insertion or a base exchange, is established in vitro in the gene of interest. The allele prepared is in turn cloned in a vector which is not replicative for *C. glutamicum* and this is then transferred
10 into the desired host of *C. glutamicum* by transformation or conjugation. After homologous recombination by means of a first "cross over" event which effects integration and a suitable second "cross-over" event which effects excision in the target gene or in the target sequence, the
15 incorporation of the mutation or of the allele is achieved. This method was used, for example, by Peters-Wendisch et al. (Microbiology 144, 915 - 927 (1998)) to eliminate the pyc gene of *C. glutamicum* by a deletion.

A deletion, insertion or a base exchange can be
20 incorporated into the metR gene or the metZ gene in this manner.

In addition, it may be advantageous for the production of L-amino acids, in particular L-methionine, to enhance, in particular to over-express, one or more enzymes of the
25 particular biosynthesis pathway, of glycolysis, of anaplerosis, of the citric acid cycle or of amino acid export, in addition to attenuation of the metR gene and/or of the metZ gene.

Thus, for example, for the preparation of L-methionine, one
30 or more genes chosen from the group consisting of

- the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),

- the *tpi* gene which codes for triose phosphate isomerase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the *pgk* gene which codes for 3-phosphoglycerate kinase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- 5 • the *pyc* gene which codes for pyruvate carboxylase (Peters-Wendisch et al. (Microbiology 144, 915 - 927 (1998)),
- the *lysC* gene which codes for a feed-back resistant aspartate kinase (Accession No.P26512),
- 10 • the *metA* gene which codes for homoserine O-acetyltransferase (ACCESSION Number AF052652),
- the *metB* gene which codes for cystathionine gamma-synthase (ACCESSION Number AF126953),
- the *aecD* gene which codes for cystathionine gamma-lyase
- 15 (ACCESSION Number M89931),
- the *glyA* gene which codes for serine hydroxymethyltransferase (JP-A-08107788),
- the *metY* gene which codes for O-acetylhomoserine sulfhydrylase (DSM 13556)
- 20 can be enhanced, in particular over-expressed.

It may furthermore be advantageous for the production of L-methionine, in addition to the attenuation of the *metR* and/or *metZ* gene, for one or more genes chosen from the group consisting of

- 25 • the *thrB* gene which codes for homoserine kinase (ACCESSION Number P08210),
- the *ilvA* gene which codes for threonine dehydratase (ACCESSION Number Q04513),

- the thrC gene which codes for threonine synthase (ACCESSION Number P23669),
- the ddh gene which codes for meso-diaminopimelate D-dehydrogenase (ACCESSION Number Y00151)
- 5 • the pck gene which codes for phosphoenol pyruvate carboxykinase (DE 199 50 409.1; DSM 13047),
- the pgi gene which codes for glucose 6-phosphate isomerase (US 09/396,478; DSM 12969),
- the poxB gene which codes for pyruvate oxidase (DE: 1995
10 1975.7)

to be attenuated, in particular for the expression thereof to be reduced.

In addition to attenuation of the metR gene and/or of the metZ gene it may furthermore be advantageous, for the
15 production of amino acids, in particular L-methionine, to eliminate undesirable side reactions (Nakayama: "Breeding of Amino Acid Producing Micro-organisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

20 The invention also provides the microorganisms prepared according to the invention, and these can be cultured continuously or discontinuously in the batch process (batch culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for the purpose of
25 production of L-amino acids, in particular L-methionine. A summary of known culture methods are described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und
30 periphere Einrichtungen (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General
5 Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981).

Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as, for example, soya oil, sunflower
10 oil, groundnut oil and coconut fat, fatty acids, such as, for example, palmitic acid, stearic acid and linoleic acid, alcohols, such as, for example, glycerol and ethanol, and organic acids, such as, for example, acetic acid, can be used as the source of carbon. These substances can be used
15 individually or as a mixture.

Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium
20 phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-
25 containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as, for example, magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and
30 vitamins, can be employed in addition to the abovementioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture.

5 Antifoams, such as, for example, fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, such as, for example, antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic
10 conditions, oxygen or oxygen-containing gas mixtures, such as, for example, air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and preferably 25°C to 40°C. Culturing is continued until a maximum of the desired product has formed. This target is
15 usually reached within 10 hours to 160 hours.

Methods for the determination of L-amino acids are known from the prior art. The analysis of L-methionine can be carried out by ion exchange chromatography with subsequent ninhydrin derivatization, as described by Spackman et al.
20 (Analytical Chemistry, 30, (1958), 1190).

The process according to the invention is used for the fermentative preparation of amino acids, in particular L-methionine.

The present invention is explained in more detail in the
25 following with the aid of embodiment examples.

The isolation of plasmid DNA from Escherichia coli and all techniques of restriction, Klenow and alkaline phosphatase treatment were carried out by the method of Sambrook et al. (Molecular Cloning. A Laboratory Manual (1989) Cold Spring
30 Harbour Laboratory Press, Cold Spring Harbor, NY, USA). Methods for transformation of Escherichia coli are also described in this handbook.

The composition of the usual nutrient media, such as LB or TY medium, can also be found in the handbook by Sambrook et al.

Example 1

- 5 Preparation of a genomic cosmid gene library from
Corynebacterium glutamicum ATCC 13032

Chromosomal DNA from Corynebacterium glutamicum ATCC 13032 was isolated as described by Tauch et al. (1995, Plasmid 33:168-179) and partly cleaved with the restriction enzyme
10 Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Code no. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Code no. 1758250). The DNA of the
15 cosmid vector SuperCos1 (Wahl et al. (1987) Proceedings of the National Academy of Sciences USA 84:2160-2164), obtained from Stratagene (La Jolla, USA, Product Description SuperCos1 Cosmid Vector Kit, Code no. 251301) was cleaved with the restriction enzyme XbaI (Amersham
20 Pharmacia, Freiburg, Germany, Product Description XbaI, Code no. 27-0948-02) and likewise dephosphorylated with shrimp alkaline phosphatase.

The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product
25 Description BamHI, Code no. 27-0868-04). The cosmid DNA treated in this manner was mixed with the treated ATCC13032 DNA and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no.27-0870-04). The ligation mixture was then
30 packed in phages with the aid of Gigapack II XL Packing Extract (Stratagene, La Jolla, USA, Product Description Gigapack II XL Packing Extract, Code no. 200217).

For infection of the E. coli strain NM554 (Raleigh et al. 1988, Nucleic Acid Research 16:1563-1575) the cells were

taken up in 10 mM MgSO₄ and mixed with an aliquot of the phage suspension. The infection and titering of the cosmid library were carried out as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 1955, Virology, 1:190) with 100 mg/l ampicillin. After incubation overnight at 37°C, recombinant individual clones were selected.

Example 2

10 Isolation and sequencing of the metR and metZ genes

The cosmid DNA of an individual colony was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme
15 Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Product No. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Product No. 1758250). After
20 separation by gel electrophoresis, the cosmid fragments in the size range of 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany).

The DNA of the sequencing vector pZero-1, obtained from
25 Invitrogen (Groningen, The Netherlands, Product Description Zero Background Cloning Kit, Product No. K2500-01) was cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid
30 fragments in the sequencing vector pZero-1 was carried out as described by Sambrook et al. (1989, Molecular Cloning: A laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was
35 then electroporated (Tauch et al. 1994, FEMS Microbiol

Letters, 123:343-7) into the E. coli strain DH5 α MCR (Grant, 1990, Proceedings of the National Academy of Sciences U.S.A., 87:4645-4649) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50 mg/l zeocin.

- 5 The plasmid preparation of the recombinant clones was carried out with Biorobot 9600 (Product No. 900200, Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain termination method of Sanger et al. (1977, Proceedings of the National Academy of Sciences U.S.A.,
- 10 74:5463-5467) with modifications according to Zimmermann et al. (1990, Nucleic Acids Research, 18:1067). The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis
- 15 of the sequencing reaction were carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1) (Product No. A124.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).
- 20 The raw sequence data obtained were then processed using the Staden program package (1986, Nucleic Acids Research, 14:217-231) version 97-0. The individual sequences of the pZero1 derivatives were assembled to a continuous contig. The computer-assisted coding region analysis was prepared
- 25 with the XNIP program (Staden, 1986, Nucleic Acids Research, 14:217-231).

- The resulting nucleotide sequence is shown in SEQ ID No. 1. Analysis of the nucleotide sequence showed two open reading frames of 567 base pairs and 1146 base pairs, which were
- 30 called the metR gene and metZ gene. The metR gene codes for a protein of 189 amino acids, the metZ gene codes for a protein of 382 amino acids.

SEQUENCE PROTOCOL

<110> Degussa-Hüls AG

5 <120> New nucleotide sequences which code for the metR and metZ genes

<130> 000369 BT

10 <140>

<141>

<160> 3

15 <170> PatentIn Ver. 2.1

<210> 1

<211> 2628

<212> DNA

20 <213> Corynebacterium glutamicum

<220>

<221> CDS

<222> (447)..(1013)

25 <223> metR gene

<220>

<221> CDS

<222> (1038)..(2183)

30 <223> metZ gene

<400> 1

```

cggtcacgtt gggatcgttg tcaaaactcc ccagtggttt cacttcataa actcgcggag 60
35 ttttccgggg aactgaaaaa catcgtccga taggccagcg tctaattcag cagcgatttc 120
   ggcagcaagc ccagcgccat taatcagagc ggtgaaataa acatgggttca tgattatgtc 180
   aggacggtaa ttagacttat gaccagggtt aaggagggtca ccagggttgaa gccgcgctat 240
40 tgttccgtgg aaaagggggc cctgatctag ctgattattc atcgcagtaa gcgctttcgg 300
   taggtgggtg aatcatcgta gtcttccgag ccccgtagacc cgatccgttt tgtgcaatcc 360
45 aatgctactc ccacagagcg ggctactttc tctaaaaatg ttctcatagt agataaaatt 420
   gttcttaaag cgacattatt gtctgc atg gaa gac gat ctc agt gct gct ctc 473
                               Met Glu Asp Asp Leu Ser Ala Ala Leu
                               1           5
50 gtc aaa gcg ctt ttc gac gcg cga acc caa cgc agg ctc tct atc tcg 521
   Val Lys Ala Leu Phe Asp Ala Arg Thr Gln Arg Arg Leu Ser Ile Ser
   10           15           20           25
55 gcg tta gct gaa tcc tcc ggt gtg tcg cga gca atg att tcc cgc gtg 569
   Ala Leu Ala Glu Ser Ser Gly Val Ser Arg Ala Met Ile Ser Arg Val
           30           35           40

```

		gaa aac gca gag gcg caa cca agc gct gca tta ctt gga cgc ctt tcc	617
		Glu Asn Ala Glu Ala Gln Pro Ser Ala Ala Leu Leu Gly Arg Leu Ser	
		45 50 55	
5		ggt gca ttg ggt atg acg ctt tcg gag ctc att gca cag gct gaa ggt	665
		Gly Ala Leu Gly Met Thr Leu Ser Glu Leu Ile Ala Gln Ala Glu Gly	
		60 65 70	
10		ggc tat gac cgg ggc gct cgg cgg tca aag cag tct gta tgg aca gat	713
		Gly Tyr Asp Arg Gly Ala Arg Arg Ser Lys Gln Ser Val Trp Thr Asp	
		75 80 85	
15		cca gct acc ggt tac aca cgg cgt gca gtg tca cag ccg tca gaa tcc	761
		Pro Ala Thr Gly Tyr Thr Arg Arg Ala Val Ser Gln Pro Ser Glu Ser	
		90 95 100 105	
20		cca cta gaa cta gtg gaa gta atg ctg cct cct ggg gcg gaa gtt ggc	809
		Pro Leu Glu Leu Val Glu Val Met Leu Pro Pro Gly Ala Glu Val Gly	
		110 115 120	
		tac cca gct gat gct tat cgt ttc atg gat cag gtg gtc tgg gta ctc	857
		Tyr Pro Ala Asp Ala Tyr Arg Phe Met Asp Gln Val Val Trp Val Leu	
		125 130 135	
25		gaa ggg gcc gtt cgt att act gaa ggt gaa gag gtc cac gaa ctt tca	905
		Glu Gly Ala Val Arg Ile Thr Glu Gly Glu Glu Val His Glu Leu Ser	
		140 145 150	
30		acg ggg gat tgt cta cgg ttt ggg cct ccg cga gat acc gac ttt gct	953
		Thr Gly Asp Cys Leu Arg Phe Gly Pro Pro Arg Asp Thr Asp Phe Ala	
		155 160 165	
35		aat ccc acc acc gta gcc act agg tat tta gtt gcc ttg gac aag cgt	1001
		Asn Pro Thr Thr Val Ala Thr Arg Tyr Leu Val Ala Leu Asp Lys Arg	
		170 175 180 185	
40		gta cct cgt gct tgatataaca agtaaggaag cctg atg aat ttt tac cca	1052
		Val Pro Arg Ala Met Asn Phe Tyr Pro	
		190	
		cca tct gta cct att aac cct gcg tgg cgt cca ccc aca gta act gtg	1100
		Pro Ser Val Pro Ile Asn Pro Ala Trp Arg Pro Pro Thr Val Thr Val	
		195 200 205 210	
45		caa gcg gga cgg cca gcc aga act cct ggt gcg ccg atg aac cca cct	1148
		Gln Ala Gly Arg Pro Ala Arg Thr Pro Gly Ala Pro Met Asn Pro Pro	
		215 220 225	
50		atc acg ttg tcc agc act tat gtt cat gat tca gaa aaa gct tat ggg	1196
		Ile Thr Leu Ser Ser Thr Tyr Val His Asp Ser Glu Lys Ala Tyr Gly	
		230 235 240	
55		cgc gat ggc aat gat gga tgg ggt gca ttt gag gct gcc atg gga act	1244
		Arg Asp Gly Asn Asp Gly Trp Gly Ala Phe Glu Ala Ala Met Gly Thr	
		245 250 255	
60		cta gat ggt ggg ttc gcg gta tct tat tct tca ggt ttg gca gcg gca	1292
		Leu Asp Gly Gly Phe Ala Val Ser Tyr Ser Ser Gly Leu Ala Ala Ala	
		260 265 270	

	acg	tcg	att	gct	gat	ttg	gtt	cct	act	ggg	ggc	aca	gtt	gtt	tta	cct	1340
	Thr	Ser	Ile	Ala	Asp	Leu	Val	Pro	Thr	Gly	Gly	Thr	Val	Val	Leu	Pro	
	275					280					285					290	
5	aaa	gct	gcc	tat	tat	ggc	gtg	acc	aat	att	ttc	gcc	agg	atg	gaa	gcc	1388
	Lys	Ala	Ala	Tyr	Tyr	Gly	Val	Thr	Asn	Ile	Phe	Ala	Arg	Met	Glu	Ala	
					295					300					305		
10	cgc	gga	agg	ctg	aag	gtt	cga	act	gtt	gat	gca	gac	aat	acc	gaa	gaa	1436
	Arg	Gly	Arg	Leu	Lys	Val	Arg	Thr	Val	Asp	Ala	Asp	Asn	Thr	Glu	Glu	
				310					315					320			
15	gtg	att	gct	gct	gct	caa	ggg	gca	gat	gtg	gtg	tgg	gtg	gaa	tcg	atc	1484
	Val	Ile	Ala	Ala	Ala	Gln	Gly	Ala	Asp	Val	Val	Trp	Val	Glu	Ser	Ile	
			325					330					335				
20	gct	aat	ccg	acg	atg	gtg	gta	gct	gat	atc	cct	gca	ata	gtc	gac	ggg	1532
	Ala	Asn	Pro	Thr	Met	Val	Val	Ala	Asp	Ile	Pro	Ala	Ile	Val	Asp	Gly	
		340					345					350					
25	gtg	cgt	ggg	ctt	gga	gtt	ttg	act	gtc	gtt	gac	gcg	act	ttc	gca	acg	1580
	Val	Arg	Gly	Leu	Gly	Val	Leu	Thr	Val	Val	Asp	Ala	Thr	Phe	Ala	Thr	
		355				360					365					370	
30	cca	ctt	cgt	caa	cgt	cca	ttg	gaa	ctt	ggg	gct	gat	att	gtg	ctt	tac	1628
	Pro	Leu	Arg	Gln	Arg	Pro	Leu	Glu	Leu	Gly	Ala	Asp	Ile	Val	Leu	Tyr	
					375					380					385		
35	tcg	gca	acc	aaa	ctt	atc	ggg	gga	cac	tct	gat	ctt	ctt	ctt	gga	gtc	1676
	Ser	Ala	Thr	Lys	Leu	Ile	Gly	Gly	His	Ser	Asp	Leu	Leu	Leu	Gly	Val	
				390					395					400			
40	gca	gtg	tgc	aag	tct	gag	cac	cat	gcg	cag	ttt	ctt	gcc	act	cac	cgt	1724
	Ala	Val	Cys	Lys	Ser	Glu	His	His	Ala	Gln	Phe	Leu	Ala	Thr	His	Arg	
			405					410					415				
45	cat	gat	cat	ggg	tca	gtg	ccg	gga	ggg	ctt	gaa	gcg	ttt	ctt	gct	ctc	1772
	His	Asp	His	Gly	Ser	Val	Pro	Gly	Gly	Leu	Glu	Ala	Phe	Leu	Ala	Leu	
		420					425					430					
50	cgt	gga	ttg	tat	tcc	ttg	gcg	gtg	cgt	ctt	gat	cga	gca	gaa	tcc	aac	1820
	Arg	Gly	Leu	Tyr	Ser	Leu	Ala	Val	Arg	Leu	Asp	Arg	Ala	Glu	Ser	Asn	
		435				440					445					450	
55	gca	gca	gaa	ctt	tcg	cgg	cga	ctt	aac	gcg	cat	cct	tcg	gtt	acc	cgc	1868
	Ala	Ala	Glu	Leu	Ser	Arg	Arg	Leu	Asn	Ala	His	Pro	Ser	Val	Thr	Arg	
					455					460					465		
50	gtc	aat	tat	cca	gga	ctt	cct	gat	gat	ccc	caa	cat	gaa	aaa	gcc	gtg	1916
	Val	Asn	Tyr	Pro	Gly	Leu	Pro	Asp	Asp	Pro	Gln	His	Glu	Lys	Ala	Val	
				470					475					480			
55	cga	gtc	cta	ccc	tct	gga	tgt	gga	aac	atg	ttg	tca	ttt	gag	ctt	gat	1964
	Arg	Val	Leu	Pro	Ser	Gly	Cys	Gly	Asn	Met	Leu	Ser	Phe	Glu	Leu	Asp	
			485					490					495				

```

gca aca cct gaa cga act gat gag att ctc gaa agc ctg tca ctt tta 2012
Ala Thr Pro Glu Arg Thr Asp Glu Ile Leu Glu Ser Leu Ser Leu Leu
500 505 510

5 acc cac gcg acc agt tgg gga ggt gtg gaa aca gcc att gaa cgt cgc 2060
Thr His Ala Thr Ser Trp Gly Gly Val Glu Thr Ala Ile Glu Arg Arg
515 520 525 530

10 acc agg cgg gat gct gaa gtg gtg gca gga gta ccg atg act ctt tgc 2108
Thr Arg Arg Asp Ala Glu Val Val Ala Gly Val Pro Met Thr Leu Cys
535 540 545

15 cgc gtt tcc gta gga att gaa gac gtt gaa gat cta tgg gaa gac ctc 2156
Arg Val Ser Val Gly Ile Glu Asp Val Glu Asp Leu Trp Glu Asp Leu
550 555 560

aac gcc tca atc gac aaa gtt ctg ggt tagaactcgt agccagtaac 2203
Asn Ala Ser Ile Asp Lys Val Leu Gly
565 570

20 cagaccttca gtgtttggtt gccactccag tgctggggcg acatgatcag cgaagttctt 2263

caggatcgac gcgttgatct caacacccat ttgggtgggg gcggtgagca tcaaggtgtc 2323

25 ggcttccatc acagctttgt cttccttgag ctggtcgatg agttcatcgg gacttccggc 2383

gtagctgca ccgaacgtgg atcgggtatc atccaggatt cctacttggg caccgccttg 2443

30 tccctgaagt ccgaaaagct cacggtcgcg gtcggtgacg atcggaaga tggacctgga 2503

gacagacaca cgtgggggtcc aatcgtgtcc ggcttctttc caagcttggc ggtagaacgc 2563

gatttgatcg gcttgcagat ccccgaagga ttggccggtg gcttcggcga cgagggtgga 2623

35 gctca 2628

<210> 2
<211> 189
40 <212> PRT
<213> Corynebacterium glutamicum

<400> 2
45 Met Glu Asp Asp Leu Ser Ala Ala Leu Val Lys Ala Leu Phe Asp Ala
1 5 10 15

Arg Thr Gln Arg Arg Leu Ser Ile Ser Ala Leu Ala Glu Ser Ser Gly
20 25 30

50 Val Ser Arg Ala Met Ile Ser Arg Val Glu Asn Ala Glu Ala Gln Pro
35 40 45

Ser Ala Ala Leu Leu Gly Arg Leu Ser Gly Ala Leu Gly Met Thr Leu
50 55 60

55 Ser Glu Leu Ile Ala Gln Ala Glu Gly Gly Tyr Asp Arg Gly Ala Arg
65 70 75 80

```

Arg Ser Lys Gln Ser Val Trp Thr Asp Pro Ala Thr Gly Tyr Thr Arg
 85 90 95
 5 Arg Ala Val Ser Gln Pro Ser Glu Ser Pro Leu Glu Leu Val Glu Val
 100 105 110
 Met Leu Pro Pro Gly Ala Glu Val Gly Tyr Pro Ala Asp Ala Tyr Arg
 115 120 125
 10 Phe Met Asp Gln Val Val Trp Val Leu Glu Gly Ala Val Arg Ile Thr
 130 135 140
 Glu Gly Glu Glu Val His Glu Leu Ser Thr Gly Asp Cys Leu Arg Phe
 15 145 150 155 160
 Gly Pro Pro Arg Asp Thr Asp Phe Ala Asn Pro Thr Thr Val Ala Thr
 165 170 175
 20 Arg Tyr Leu Val Ala Leu Asp Lys Arg Val Pro Arg Ala
 180 185

 <210> 3
 25 <211> 382
 <212> PRT
 <213> Corynebacterium glutamicum

 <400> 3
 30 Met Asn Phe Tyr Pro Pro Ser Val Pro Ile Asn Pro Ala Trp Arg Pro
 1 5 10 15
 Pro Thr Val Thr Val Gln Ala Gly Arg Pro Ala Arg Thr Pro Gly Ala
 20 25 30
 35 Pro Met Asn Pro Pro Ile Thr Leu Ser Ser Thr Tyr Val His Asp Ser
 35 40 45
 Glu Lys Ala Tyr Gly Arg Asp Gly Asn Asp Gly Trp Gly Ala Phe Glu
 40 50 55 60
 Ala Ala Met Gly Thr Leu Asp Gly Gly Phe Ala Val Ser Tyr Ser Ser
 65 70 75 80
 45 Gly Leu Ala Ala Ala Thr Ser Ile Ala Asp Leu Val Pro Thr Gly Gly
 85 90 95
 Thr Val Val Leu Pro Lys Ala Ala Tyr Tyr Gly Val Thr Asn Ile Phe
 100 105 110
 50 Ala Arg Met Glu Ala Arg Gly Arg Leu Lys Val Arg Thr Val Asp Ala
 115 120 125
 Asp Asn Thr Glu Glu Val Ile Ala Ala Ala Gln Gly Ala Asp Val Val
 55 130 135 140
 Trp Val Glu Ser Ile Ala Asn Pro Thr Met Val Val Ala Asp Ile Pro
 145 150 155 160
 60 Ala Ile Val Asp Gly Val Arg Gly Leu Gly Val Leu Thr Val Val Asp

26

	165					170					175					
	Ala	Thr	Phe	Ala	Thr	Pro	Leu	Arg	Gln	Arg	Pro	Leu	Glu	Leu	Gly	Ala
				180					185					190		
5	Asp	Ile	Val	Leu	Tyr	Ser	Ala	Thr	Lys	Leu	Ile	Gly	Gly	His	Ser	Asp
			195					200					205			
10	Leu	Leu	Leu	Gly	Val	Ala	Val	Cys	Lys	Ser	Glu	His	His	Ala	Gln	Phe
		210					215					220				
	Leu	Ala	Thr	His	Arg	His	Asp	His	Gly	Ser	Val	Pro	Gly	Gly	Leu	Glu
	225					230					235					240
15	Ala	Phe	Leu	Ala	Leu	Arg	Gly	Leu	Tyr	Ser	Leu	Ala	Val	Arg	Leu	Asp
					245					250					255	
	Arg	Ala	Glu	Ser	Asn	Ala	Ala	Glu	Leu	Ser	Arg	Arg	Leu	Asn	Ala	His
				260					265					270		
20	Pro	Ser	Val	Thr	Arg	Val	Asn	Tyr	Pro	Gly	Leu	Pro	Asp	Asp	Pro	Gln
			275					280					285			
	His	Glu	Lys	Ala	Val	Arg	Val	Leu	Pro	Ser	Gly	Cys	Gly	Asn	Met	Leu
25		290					295					300				
	Ser	Phe	Glu	Leu	Asp	Ala	Thr	Pro	Glu	Arg	Thr	Asp	Glu	Ile	Leu	Glu
	305					310					315					320
30	Ser	Leu	Ser	Leu	Leu	Thr	His	Ala	Thr	Ser	Trp	Gly	Gly	Val	Glu	Thr
					325					330					335	
	Ala	Ile	Glu	Arg	Arg	Thr	Arg	Arg	Asp	Ala	Glu	Val	Val	Ala	Gly	Val
				340					345					350		
35	Pro	Met	Thr	Leu	Cys	Arg	Val	Ser	Val	Gly	Ile	Glu	Asp	Val	Glu	Asp
			355					360					365			
	Leu	Trp	Glu	Asp	Leu	Asn	Ala	Ser	Ile	Asp	Lys	Val	Leu	Gly		
40		370					375					380				

45

Patent claims

1. An isolated polynucleotide from coryneform bacteria,
which comprises a polynucleotide sequence which codes
for the metR and/or metZ genes, chosen from the group
consisting of
 - a) polynucleotide which is identical to the extent of
at least 70% to a polynucleotide which codes for a
polypeptide which comprises the amino acid
sequence of SEQ ID No. 2,
 - b) polynucleotide which is identical to the extent of
at least 70% to a polynucleotide which codes for a
polypeptide which comprises the amino acid
sequence of SEQ ID No. 3,
 - c) polynucleotide which codes for a polypeptide which
comprises an amino acid sequence which is
identical to the extent of at least 70% to the
amino acid sequence of SEQ ID No. 2,
 - d) polynucleotide which codes for a polypeptide which
comprises an amino acid sequence which is
identical to the extent of at least 70% to the
amino acid sequence of SEQ ID No. 3,
 - e) polynucleotide which is complementary to the
polynucleotides of a), b), c) or d), and
 - f) polynucleotide comprising at least 15 successive
nucleotides of the polynucleotide sequence of a),
b), c), d) or e).
2. A polynucleotide as claimed in claim 1, wherein the
polynucleotide is a preferably recombinant DNA which
is capable of replication in coryneform bacteria.
3. A polynucleotide as claimed in claim 1, wherein the
polynucleotide is an RNA.

4. A polynucleotide as claimed in claim 2, comprising the nucleic acid sequence as shown in SEQ ID No. 1.
5. A DNA as claimed in claim 2 which is capable of replication, comprising
 - 5 (i) the nucleotide sequence shown in SEQ ID No. 1, or
 - (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
 - 10 (iii) at least one sequence which hybridizes with the sequence complementary to sequence (i) or (ii), and optionally
 - (iv) sense mutations of neutral function in (i).
6. A DNA as claimed in claim 5 which is capable of replication, wherein the hybridization of sequence
 - 15 (iii) is carried out under a stringency corresponding to at most 2x SSC.
7. A polynucleotide sequence as claimed in claim 1, which codes for a polypeptide which comprises the amino acid sequence in SEQ ID No. 2 and/or SEQ ID No. 3.
- 20 8. A coryneform bacterium in which the metR gene and/or metZ gene is or are attenuated, in particular eliminated.
9. A process for the fermentative preparation of L-amino acids, in particular L-methionine, which comprises
 - 25 carrying out the following steps:
 - a) fermentation of the coryneform bacteria which produce the desired L-amino acid and in which at least the metR and/or metZ gene or nucleotide sequences which code for them are attenuated;

- b) concentration of the L-amino acid in the medium or in the cells of the bacteria, and
 - c) isolation of the L-amino acid.
10. A process as claimed in claim 9, wherein bacteria in which further genes of the biosynthesis pathway of the desired L-amino acid are additionally enhanced are employed.
11. A process as claimed in claim 9, wherein bacteria in which the metabolic pathways which reduce the formation of the desired amino acid are at least partly eliminated are employed.
12. A process as claimed in claim 9, wherein the expression of the polynucleotide(s) which code(s) for the metR and/or the metZ gene is reduced or attenuated.
13. A process as claimed in claim 9, wherein the catalytic properties of the polypeptides (enzyme protein) for which the polynucleotides metR and/or metZ code are attenuated.
14. A process as claimed in claim 9, wherein for the preparation of L-amino acids, in particular L-methionine, coryneform microorganisms in which at the same time one or more of the genes chosen from the group consisting of
- 14.1 the lysC gene which codes for a feed back resistant aspartate kinase,
 - 14.2 the gap gene which codes for glycerolaldehyde 3-phosphate dehydrogenase,
 - 14.3 the pyc' gene which codes for pyruvate carboxylase,

- 14.4 the tpi gene which codes for triose phosphate isomerase
- 14.5 the metA gene which codes for homoserine O-acetyltransferase
- 5 14.6 the metB gene which codes for cystathionine gamma-synthase
- 14.7 the pgk gene which codes for 3-phosphoglycerate kinase
- 10 14.8 the aecD gene which codes for cystathionine gamma-lyase
- 14.9 the glyA gene which codes for serine hydroxymethyltransferase
- 14.10 the metY gene which codes for O-acetylhomoserine sulfhydrylase
- 15 is or are enhanced or over-expressed are fermented.
- 15. A process as claimed in claim 9, wherein for the preparation of L-amino acids, in particular L-methionine, coryneform microorganisms in which at the same time one or more of the genes chosen from the group consisting of
 - 15.1 the thrB gene which codes for homoserine kinase
 - 15.2 the ilvA gene which codes for threonine dehydratase
 - 25 15.3 the thrC gene which codes for threonine synthase
 - 15.4 the ddh gene which codes for meso-diaminopimelate D-dehydrogenase

- 15.5 the pck gene which codes for phosphoenol
pyruvate carboxykinase
- 15.6 the pgi gene which codes for glucose 6-
phosphate isomerase
- 5 15.7 the poxB gene which codes for pyruvate oxidase
is or are attenuated are fermented.
- 16. A coryneform bacterium which contains a vector which
carries a polynucleotide as claimed in claim 1f.
- 17. A process as claimed in one or more of the preceding
10 claims, wherein microorganisms of the species
Corynebacterium glutamicum are employed.
- 18. A process for discovering RNA, cDNA and DNA in order
to isolate nucleic acids, or polynucleotides or genes
which code for O-succinylhomoserine sulfhydrylase
15 (metZ) and/or the transcription activator MetR or have
a high similarity with the sequence of the O-
succinylhomoserine sulfhydrylase (metZ) gene or of the
transcription activator MetR, which comprises
employing the polynucleotide comprising the
20 polynucleotide sequences as claimed in claims 1, 2, 3
or 4 as hybridization probes.

Abstract

The invention relates to polynucleotides from coryneform bacteria which code for the metR and/or metZ genes and comprise polynucleotide sequences, chosen from the group
5 consisting of

- a) polynucleotide which is identical to the extent of at least 70% to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 2,
- 10 b) polynucleotide which is identical to the extent of at least 70% to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 3,
- c) polynucleotide which codes for a polypeptide which
15 comprises an amino acid sequence which is identical to the extent of at least 70% to the amino acid sequence of SEQ ID No. 2,
- d) polynucleotide which codes for a polypeptide which comprises an amino acid sequence which is
20 identical to the extent of at least 70% to the amino acid sequence of SEQ ID No. 3,
- e) polynucleotide which is complementary to the polynucleotides of a), b), c) or d), and
- f) polynucleotide comprising at least 15 successive
25 nucleotides of the polynucleotide sequences of a), b), c), d) or e)

and a process for the fermentative preparation of L-amino acids using coryneform bacteria in which at least the metR gene and/or the metZ gene is present in attenuated form,
30 and the use of polynucleotides which comprise the sequences according to the invention as hybridization probes.